

Cerebrolysin Dosing Chart

A mixture of peptides derived from pig brain, used in some countries for stroke and dementia. Not approved in the US; evidence is contested. Commonly discussed doses: 10 mg–60 mg. Dosing in countries where it is used is by milliliters of the preparation, given by injection or IV over courses. Figures shown are total milligrams in micrograms. It is administered clinically, not as a home reconstitution.

thelastpeptide.com · Medically reviewed by Charles Maddix, MSN, APRN, PMHNP-BC · Last reviewed 2026-08-02

FDA STATUS

Not FDA-approved in the United States. Cerebrolysin is used in some other countries for stroke and dementia, but its evidence base is contested and it is not FDA-approved.

10 mg vial — units to draw (U-100 syringe)

Water added	mg/mL	10 mg	30 mg	60 mg
1 mL	10	100 u	—	—
2 mL	5	—	—	—
3 mL	3.33	—	—	—

30 mg vial — units to draw (U-100 syringe)

Water added	mg/mL	10 mg	30 mg	60 mg
1 mL	30	33.3 u	100 u	—
2 mL	15	66.7 u	—	—
3 mL	10	100 u	—	—

"—" means the draw exceeds one syringe at that concentration. Interactive calculator for any combination: thelastpeptide.com/peptides/cerebrolysin

Cerebrolysin dose figures are commonly discussed reference information with no approved labeling unless stated above. The Last Peptide is an educational reference and sells nothing. Dosing figures shown here are commonly discussed reference information, not prescriptions or medical advice. Many peptides listed are not FDA-approved. Talk to a licensed clinician before using any peptide.