

MK-677 Dosing Chart

An oral growth-hormone secretagogue (not a peptide) studied in human trials. Investigational; raises appetite and blood sugar. Commonly discussed doses: 10 mg–50 mg. MK-677 is taken orally in milligrams, not reconstituted or injected. Figures shown are total milligrams in micrograms — 25,000 mcg is 25 mg. It is a capsule/liquid oral drug, not a reconstituted peptide.

thelastpeptide.com · Medically reviewed by Charles Maddix, MSN, APRN, PMHNP-BC · Last reviewed 2026-08-02

FDA STATUS

Not FDA-approved for any use. MK-677 (ibutamoren) is an investigational oral drug that has been through human trials but is not approved. It is prohibited in sport.

10 mg vial — units to draw (U-100 syringe)

Water added	mg/mL	10 mg	25 mg	50 mg
1 mL	10	100 u	—	—
2 mL	5	—	—	—
3 mL	3.33	—	—	—

25 mg vial — units to draw (U-100 syringe)

Water added	mg/mL	10 mg	25 mg	50 mg
1 mL	25	40 u	100 u	—
2 mL	12.5	80 u	—	—
3 mL	8.33	—	—	—

"—" means the draw exceeds one syringe at that concentration. Interactive calculator for any combination: thelastpeptide.com/peptides/mk677

MK-677 dose figures are commonly discussed reference information with no approved labeling unless stated above. The Last Peptide is an educational reference and sells nothing. Dosing figures shown here are commonly discussed reference information, not prescriptions or medical advice. Many peptides listed are not FDA-approved. Talk to a licensed clinician before using any peptide.